

Support for Clients using GLP-1 Agonists

PBMT and WBV is suitable to support clients who are using GLP-1 medications.

Photobiomodulation supports metabolic health and is not a substitute for appropriate medical care.

Side Effects Related to GLP-1 Agonists

- Loss of Muscle Mass
- Decreased bone density
- Rapid loss of facial fat that outpaces collagen production leading to prominent wrinkles and hollow cheeks
- Optic neuropathy
- Pancreatitis (inflammation)
- Dehydration and kidney damage
- Rebound weight gain after discontinuation

Support for Clients using GLP-1 Agonists

Concerns that can be addressed with PBMT and WBV

- PBM and WBV promotes maintenance of Muscle Mass
- WBV promotes maintenance of bone density
- PBM stimulates skin regeneration, collagen and elastin production and leading to improved skin density
- PBM activates anti-inflammatory signaling pathways,
- PBM supports organs that require large amounts of cellular energy for optimum functioning like the eyes, nerves, muscles and kidneys.
- PBM and WBV support metabolic health and healthy fat loss, while building muscle.
- PBM and WBV can decrease the rebound weight gain experienced after stopping GLP-1 medications.

A Point to note: Red and Infrared Light have a potent partnership with UV Light.

Red and Infrared light are regenerative and supportive of metabolism, BUT UV light is a natural GLP-1 regulator, decreasing appetite and increasing metabolism. Vitamin D deficiency is a key obstacle to metabolic health. Encourage your clients to build a robust vitamin D level built from Ultraviolet light exposure.

Your clients will do better if they also re-establish their circadian rhythms by exposure to early morning sunlight, bright sunlight during the day (take light breaks) and shift to dimmed lighting (warm) at night.

Support for Clients using GLP-1 Agonists

DO NOT TREAT

- Any active systemic infection (fevers)
- Autoimmune condition in a flare-up, or client on high dose steroids, clients using high dose cortisone/steroids (only treat once tapering to a low dose has begun)

Note any medications being used concurrently.

PROTOCOL

10 minutes PBM on back

10 min PBM on front

5 minutes of WBV before and after the PBM session (if applicable)

20-30cm Panel at 20-30cm depending on heat tolerance

- **Eye protection mandatory**

Be aware that clients may have low blood glucose levels and experience fatigue earlier on the vibration plate due to decreased appetite and muscle loss.